

Deep Venous Thrombosis After Orthopedic Surgery in Adult Cancer Patients

PATRICK P. LIN, MD,¹ DENNIS GRAHAM, RN, NP,² LUCY E. HANN, MD,³
PATRICK J. BOLAND, MD,¹ AND JOHN H. HEALEY, MD^{1*}

¹Orthopaedic Surgery Service, Memorial Sloan-Kettering Cancer Center,
New York, New York

²Department of Surgery, Memorial Sloan-Kettering Cancer Center, New York, New York

³Department of Radiology, Memorial Sloan-Kettering Cancer Center, New York, New York

Background and Objectives: Patients with cancer and patients undergoing major orthopedic procedures are two groups at risk of deep venous thrombosis (DVT). The objective was to determine the rate of venous thromboembolic disease in patients with a malignant neoplasm and major orthopaedic surgery of the lower limb.

Methods: The study included 169 patients. All patients were given knee-high intermittent pneumatic compression devices for prophylaxis. Postoperative surveillance for thrombosis was performed on all patients with venous duplex doppler ultrasonography.

Results: Proximal DVT occurred in 24 of 169 patients (14.2%). One patient (0.6%) developed a symptomatic, nonfatal pulmonary embolus (PE). The development of DVT was not associated with age, sex, type of surgery, type of neoplasm, location, or pathologic fracture. The addition of anticoagulant medication such as warfarin did not significantly reduce the rate of DVT in a subset of 54 patients. In three patients, the DVT occurred only in the contralateral limb, and in four patients, there were bilateral DVTs.

Conclusions: When intermittent compression boots were used for prophylaxis in conjunction with ultrasound screening, the risk of proximal DVT was substantial (14.2%), but the rate of symptomatic PE was low (0.6%). *J. Surg. Oncol.* 1998;68:41–47. © 1998 Wiley-Liss, Inc.

KEY WORDS: thromboembolism; pulmonary embolus; postoperative complications; duplex doppler ultrasonography; orthopedics

INTRODUCTION

It has been generally recognized that patients with cancer are at increased risk for thromboembolic disease [1–6]. This observation has been made for many different types of tumors and malignancies, especially those involving the pelvis [7–15]. It is somewhat surprising that the problem has received scant attention in the orthopedic literature, especially in light of the strong association between various orthopedic conditions and venous thrombosis. Patients who undergo total hip and total knee replacements are at high risk for developing thromboembolic complications [16–18]. Without prophylaxis, the overall rate of thromboembolic events has been estimated to be 45–69% for total hip replacement [16,19–23]

and 52–84% for total knee replacement [24–29]. The rate of pulmonary emboli has been reported to be 3.2–23% for hip replacement [16,17,30–33] and 2.7–24% for knee replacement [32–35]. Patients who have hip, pelvic, and acetabular fractures are also notably at high risk for thromboembolic disease with overall rates of deep ve-

Investigation performed at Memorial Sloan-Kettering Cancer Center, affiliated with Cornell University Medical College, New York, New York.

Grant sponsor: New York Marathon Limb Preservation Fund.

*Correspondence to: John H. Healey, MD, Orthopaedic Surgery Service, Department of Surgery, Memorial Sloan-Kettering Cancer Center, 1275 York Ave., New York, NY 10021. Telephone No.: (212) 639-7610; Fax No.: (212) 794-4015. E-mail: healeyj@mskcc.org

Accepted 17 February 1998

nous thrombosis ranging from 39–75% [21,36–42]. It has thus been assumed that orthopedic patients with malignant disease would be at the highest risk of thromboembolism [5].

Patients with musculoskeletal tumors have the potential for fulfilling Virchow's classic triad. (1) The neoplastic process may liberate factors that affect the clotting process and induce hypercoagulability [1,43,44]. (2) Circulatory stasis is induced by bulky tumors, decreased mobility, long operation times, and unnatural positioning of limbs during surgery. (3) Pathologic fractures and surgical trauma can each lead to endothelial damage.

Although it seems plausible that patients with neoplasms undergoing orthopedic surgery would be at increased risk for deep venous thrombosis, there has been no study that attempts to quantify the magnitude of the problem or assess the adequacy of current methods of prophylaxis. The lack of published data probably stems from the numerous and substantial difficulties in studying this problem. Primary musculoskeletal tumors are rare, and it is not easy to obtain sufficient numbers of patients for a meaningful study. The population of patients with orthopedic neoplasms is heterogeneous, and there is a wide variety of lesions, locations, procedures, ages, and other patient characteristics involved. Finally, there have been no previous studies on the natural history of DVT in these patients, which precludes the possibility of using historical controls to measure the effectiveness of current methods of prophylaxis.

The goal of the present study is to determine the prevalence of DVT in patients with neoplasms who undergo orthopedic surgery and receive basic prophylaxis in the form of pneumatic compression stockings. The study focuses on the patients who seem most at risk for developing thrombosis—those with active malignant neoplasms who undergo major procedures on the pelvis or lower limbs. Specific characteristics of patients are examined to determine whether certain subsets are at higher risk of thrombosis.

MATERIALS AND METHODS

Study Design

A prospective study was undertaken to evaluate the rate of DVT in cancer patients undergoing major musculoskeletal surgery of the lower limbs. All 169 patients in the study were treated at this institution between June 1994 and August 1996. A protocol for DVT prophylaxis and screening was established at the beginning of the study (see below).

Patients

Patients had to satisfy the following criteria for inclusion in the study. Age was 21 years or older. Major musculoskeletal surgery was performed on the pelvis or lower limbs. The procedure had to result in significantly diminished mobility and require hospitalization of at

least 4 days. An active malignant neoplasm had to be present at the time of surgery. Patients who had a previous history of cancer but no detectable disease at the time of surgery were not included.

Patients were excluded for the following reasons: (1) minor operations, such as simple biopsies, (2) a previously placed inferior vena caval filter, (3) diagnosis of a DVT prior to surgery, (4) no detectable malignant disease at the time of operation, (5) an impediment to obtaining a venous ultrasound scan, such as brachtherapy catheters or a cast, and (6) amputation. Patients who underwent amputations were excluded since they were expected to have thrombosis at the level of the divided vein.

Of the 193 patients who met eligibility requirements, 169 patients were included in the study. The remaining 24 patients were not included because they did not have a postoperative ultrasound scan. Ninety-five females and 74 males were entered into the study. Ages ranged from 21 yr to 92 yr with a mean of 59.7 yr. A total of 103 patients had metastatic disease to the surgical site; 50 patients had primary or locally recurrent tumors; and 16 patients had concurrent neoplasms that did not directly involve the surgical site.

Protocol for DVT Prophylaxis and Management

Prior to surgery, all patients were examined carefully for signs of venous thromboembolism. Seventeen patients were studied with pre-operative duplex doppler ultrasonography, and six patients were found to have a DVT. These patients were treated with inferior vena caval filters and excluded from the present analysis.

Patients were given knee-high pneumatic intermittent compression devices (7 Venodyne, Advanced Instruments, Norwood, MA) during and after surgery. The device inflated to 30–40 mm Hg for 12 sec, then deflated for 48 sec, thus cycling once every minute. During surgery, the sleeve was placed only on the uninvolved extremity. After surgery, the sleeves were worn constantly, and they were placed on both extremities whenever feasible. Intermittent compression was continued until the ultrasound scan results were known and the patient ambulatory. All data are based upon intent to treat with intermittent compression boots and not on strict compliance with wearing the boots.

Fifty-four patients received anticoagulant medications in addition to intermittent compression devices postoperatively. The decision to give the medications was made on a case-by-case basis at the discretion of the attending physician. The administration of all anticoagulant medications was carefully monitored, and complete data on these medications was verified against hospital pharmacy computer records.

All patients were screened for postoperative DVT (see below). Patients who were diagnosed with venous throm-

boembolism were treated acutely with an intravenous heparin infusion. Patients were subsequently managed with either an inferior vena caval filter and/or oral warfarin for a total of 3 mo.

Screening for Venous Thromboembolism

Duplex ultrasound scans were performed on all patients between the fourth and seventh postoperative day. The timing of the scans varied since they were not routinely available on weekends. Pulmonary ventilation-perfusion scans were performed only on symptomatic patients suspected of having a pulmonary embolism. Routine surveillance lung scans were not performed.

According to American College of Radiology protocol standards [45], ultrasonography was performed from the saphenofemoral junction to the popliteal trifurcation to include the common femoral vein, origin of the deep femoral vein, superficial femoral vein, and popliteal vein. Calf veins were not routinely evaluated. Studies were performed with standard compression and doppler technique as described by Cronan et al. [46] and all studies were monitored by an experienced radiologist. Venous thrombosis was diagnosed when veins were noncompressible. Only DVTs in the femoral-popliteal system were scored as positive. Thromboses of superficial veins, such as the saphenous vein, and calf veins were not counted as DVTs.

Statistics

All data were entered into an Excel (8 Microsoft Corporation) database and then transferred to SPSS (8 SPSS) for statistical analysis. The Chi-square test was used to compare the various groups with respect to occurrence of DVT. Fisher's exact test was used for small numbers (i.e., when the expected frequency of a cell in a 2×2 table was <5). The student's *t*-test was used to compare the means of the ages of patients with and without DVT. Statistical significance was defined as $P < 0.05$. Correlative studies were performed separately upon patients who received supplemental chemical anticoagulation and those who did not. Since there were no differences between the groups in any of the demographic or treatment factors analyzed, the analysis is presented for the entire study population.

RESULTS

Of the 169 patients who underwent postoperative venous ultrasonography, 22 patients had a positive scan for a deep proximal DVT in the immediate postoperative period between the fourth and seventh day. Two additional patients had negative scans initially but developed symptomatic DVTs subsequently, one at 1 mo and one at 3 mo. Both of these DVTs were considered to be postoperative DVTs. Thus a total of 24/169 patients (14.2%) were found to have a proximal deep DVT. Three patients

TABLE I. Postoperative Deep Venous Thrombosis (DVT) in Adults With Cancer Undergoing Orthopedic Surgery: Location of Surgery and Occurrence of DVT

Location	No. of patients	No. of patients with DVT	% DVT
Pelvis and sacrum	30	5	16.7
Proximal femur and hip	96	11	11.5
Mid femur and thigh	19	3	15.8
Distal femur and knee	16	4	25.0
Tibia, fibula, and calf	8	1	12.5

had a DVT isolated to the contralateral limb, and four patients had bilateral DVTs, making a total of 7/24 patients (29% of the patients with DVTs) who had a remote thrombus away from the surgical side.

Most patients with DVTs were treated initially with an intravenous heparin infusion. An inferior vena caval filter was placed without complication in nine patients. The other patients were treated with oral warfarin. One of the patients on warfarin developed a symptomatic PE on the sixth postoperative day. The lung scan showed a ventilation-perfusion mismatch, which was interpreted as highly probable for a PE. This patient continues to be alive 2 yr and 1 mo after surgery. The overall prevalence of symptomatic, perioperative PE in this series was 0.6%.

The occurrence of a DVT was not associated with gender. Twelve of 74 males (16.2%) and 12 of 95 females (12.6%) developed a DVT ($P = 0.52$, two-tailed test). The presence of a fracture did not significantly increase the risk of a DVT. Eight of 60 patients (13.3%) with a fracture had a DVT, whereas 16 of 109 patients (14.7%) without a fracture developed a DVT ($P = 0.81$, two-tailed test). No single location of surgery was associated with a significantly increased or decreased rate of DVT (Table I).

Age was not associated with the presence of a DVT ($P = 0.48$, two-tailed test). The mean age of patients with DVTs was 61.5 ± 12.3 standard deviation (S.D.) yr, and the mean age of patients without DVTs was 59.4 ± 13.4 yr. The youngest patient to have a DVT was 38 yr old. None of the 12 patients between 21 and 38 had a DVT, but this was not statistically significant ($P = 0.15$, one-tailed test).

The most common operative procedures are listed in Table II. The risk of postoperative DVT was not significantly increased or decreased for any single procedure. For total knee replacements, which had the highest rate of DVTs, 4 of 16 patients (25%) developed DVTs ($P = 0.25$, two-tailed test).

The most common pathologic diagnoses are listed in Table III. Again, the risk of DVT was not significantly increased or decreased for a particular diagnosis. Chondrosarcoma was associated with the highest rate of DVT (5 of 18 patients [27.8%]), but this was not statistically significant ($P = 0.14$, two-tailed test). Although none of

TABLE II. Postoperative Deep Venous Thrombosis (DVT) in Adults With Cancer Undergoing Orthopedic Surgery: Rate of CVT for Different Procedures

Procedure	No. of patients	No. of patients with DVT	% with DVT
Internal hemipelvectomy	11	2	18.2
Total hip replacement	23	2	8.7
Hip hemiarthroplasty	83	10	12.0
Total knee replacement	16	4	25.0
Soft tissue	15	3	20.0
All other procedures	21	3	14.3

TABLE III. Postoperative Deep Venous Thrombosis (DVT) in Adults With Cancer Undergoing Orthopedic Surgery: Effect of Pathologic Diagnosis

Diagnosis	No. of patients	No. of patients with DVT	% with DVT
Breast	35	5	14.3
Sarcoma ^a	32	5	15.6
Chondrosarcoma	18	5	27.8
Lung	16	3	18.8
Prostate	16	1	6.3
Renal	13	3	23.1
Multiple myeloma	11	0	0
All other	28	2	7.1

^aOther than chondrosarcoma.

the 11 patients with multiple myeloma had a DVT, this also was not statistically significant ($P = 0.37$, two-tailed test).

Patients were analyzed from the perspective of whether they had primary, metastatic, or concurrent lesions (Table IV). Concurrent lesions included cases such as nonpathologic fractures in patients with cancer. Locally recurrent tumors were combined with primary tumors. There was not a statistically significant relationship between these categories and the occurrence of a DVT ($P = 0.37$, two-tailed test).

A total of 54 patients received some form of anticoagulation postoperatively (Table V). Seventeen patients received low dose warfarin, and 32 received subcutaneous heparin. One patient received low-molecular dextran for a vascular repair, and four patients received aspirin. The administration of anticoagulant medication was not associated with a significantly reduced risk of DVT when compared to patients who received only pneumatic intermittent compression. Postoperative DVTs occurred in 7 of 54 patients (13.0%) who received both intermittent compression and anticoagulant medication and 17 of 115 patients (14.8%) who received only pneumatic intermittent compression ($P = 0.48$, one-tailed test). Only one of 17 patients (5.9%) who received warfarin and intermittent compression developed a DVT, but this was not significantly different from patients who received intermittent compression only ($P = 0.29$, one-tailed test).

TABLE IV. Postoperative Deep Venous Thrombosis (DVT) in Adults With Cancer Undergoing Orthopedic Surgery: Effect of Type of Lesion

Type of lesion	No. of patients	No. of patients with DVT	% with DVT
Primary or recurrent ^a	50	10	20.0
Concurrent	16	2	12.5
Metastatic	103	12	11.7

^aLocally recurrent in the same site as original primary tumor.

TABLE V. Postoperative Deep Venous Thrombosis (DVT) in Adults With Cancer Undergoing Orthopedic Surgery: Effect of Anticoagulant Medications

Medication	No. of patients	No. of patients with DVT	% with DVT
None	115	17	14.8
Warfarin	17	1	5.9
Heparin	32	5	15.6
All other	5	1	20.0

Thirty-five patients died during the period of follow-up. Six of these patients were known to have a DVT. There was not a statistically significant relationship between death and the occurrence of a DVT ($P = 0.37$, one-tailed test). Of the 13 patients who died within 3 mo of surgery, one was known to have a DVT, and this was not statistically significant ($P = 0.42$, one-tailed test).

DISCUSSION

Little information is available on DVTs in cancer patients having orthopedic surgery despite the widely held notion that such patients would be at increased risk of venous thromboembolism. This is a difficult subject to study since orthopedic oncology encompasses the treatment of many diverse and rare disorders. Consequently, there have been no studies on the natural history of DVTs in this patient population, and it is not known what the rate of DVT would be without prophylaxis. It is also not known what form of prophylaxis would be most appropriate for these patients. Although anticoagulant medications are widely used in orthopedic patients, the medications may not be ideal for debilitated patients who sustain considerable blood loss during surgery.

The goal of the present study was to determine the rate of proximal DVT with pneumatic compression boots. This form of prophylaxis has been shown to have some efficacy with essentially no risk of serious complications [26,34,47–51]. The devices are widely used and applicable to nearly all patients. Even in cases where postoperative dressings and splints interfere with the device, there has been reported beneficial systemic effect from placing the device on an uninvolved extremity [52–54].

The study would have benefitted from inclusion of a placebo control to assess the efficacy of the pneumatic

compression boots, but such a study design is difficult to justify from both an ethical and legal standpoint because of the potential catastrophic outcome of thromboembolism and the prevailing concept that cancer patients are at high risk for thromboembolic disease [1–6]. Current practice guidelines by the state Physician Review Organization dictate that all hip and knee replacement and other high-risk patients must receive prophylaxis against DVT.

The prevalence of proximal DVT in this series was 14.8% (17/115) for compression boots alone. The figure is relatively high when compared to other orthopedic series employing external pneumatic compression, which have reported rates of proximal DVT ranging from 6–15% [49–51,55–58]. Since ultrasound was used for diagnosis, the rate of DVT reported here is probably a conservative estimate, and the true rate is likely to be even higher. Duplex doppler ultrasound has certain advantages, including noninvasiveness, portability, and relatively low cost [46,59–62]. The specificity is excellent and approaches 97–100% for proximal DVTs [63–66]. However, a recent meta-analysis has shown that the overall sensitivity in well-controlled studies is only 79% for postoperative orthopedic patients [65]. It has also been observed that the accuracy of the test depends on the experience and skill of the operator [67].

Despite the substantial rate of DVT with compression boots alone, the rate of symptomatic PE was low (0.6%) in this series. The paucity of PEs may be attributable in part to the screening protocol for DVT, which included a preoperative clinical evaluation, doppler scans on symptomatic preoperative patients, and surveillance doppler scans on all postoperative patients. Whereas boots may be only moderately effective in this patient population, careful screening may help improve the ultimate outcome by reducing the risk of major, symptomatic PE.

The occurrence of subclinical, small PEs may have escaped detection since only symptomatic patients were studied with lung scans. However, there was no indication that undetected thromboembolic disease had an impact on outcome, and a statistically significant relationship could not be found between the occurrence of DVT and postoperative death. Of the 13 patients who died during the first 3 mo after surgery, only one was known to have had a DVT. Nevertheless, since autopsies were not routinely performed, it is not certain whether thromboembolism contributed to these patients' death.

Anticoagulant medications are effective prophylactic agents, but are associated with hemorrhagic and other complications [30,68–76]. Patients with tumors often have abnormalities of coagulation that promote bleeding and consumption of clotting factors [77,78]. Due to high blood loss and the frequent necessity to operate through highly vascular tissue planes, the patients in this study were expected to have a greater risk of postoperative

bleeding and wound complications than nononcologic orthopedic patients. Consequently, most patients in this study were managed with compression boots alone.

Fifty-four patients received anticoagulation postoperatively. No significant difference in the rate of DVT was found between patients that received anticoagulant medications and those that had pneumatic compression devices only. The data can be interpreted several ways. Chemical anticoagulation may have added no protection against DVT beyond that of intermittent compression alone. Alternatively, there may have been selection bias in the choice of patients to receive chemical prophylaxis, and this may have masked the effect of anticoagulation. Finally, there may have been insufficient numbers of patients to detect a significant difference. Clearly, more work is needed to define further the role of chemical prophylaxis for these patients.

The analysis demonstrated that no particular diagnosis, procedure, or characteristic was associated with a significantly higher or lower rate of DVT. Therefore, all cancer patients having major orthopedic surgery of the pelvis or lower extremities should be considered to be at substantial risk of thromboembolic disease, and a high index of suspicion should be maintained for all patients. It is possible that with a larger sample size, certain subsets of patients eventually may be found to have greater or less risk of DVT, and this may have an impact upon the choice of prophylaxis for that particular group of patients.

An interesting observation was that three patients had DVTs isolated to the contralateral limb and four additional patients had bilateral DVTs. Thus a total of seven of 24 patients (29%) had a proximal thrombus away from the site of surgery. The finding supports the notion that in addition to local, mechanical factors that promote the formation of DVT, there are also systemic factors that stimulate thrombosis in cancer patients. The finding supports the use of bilateral ultrasound examinations for screening as opposed to unilateral scans in this patient population.

CONCLUSIONS

The overall prevalence of proximal, postoperative DVT was 14.8% with compression boots alone. It is not known which cancer patients undergoing major lower extremity orthopedic procedures may require chemical prophylaxis for DVT. Any decision must be balanced against the hemorrhagic and general risks in this debilitated population. There was no clinical subset of patients that was at significantly less or greater risk than other patients, and thus a high index of suspicion should be maintained for all patients. When compression boots were used in conjunction with a screening program that included routine surveillance postoperative ultrasound scans, the rate of symptomatic PE was 0.6%.

ACKNOWLEDGMENTS

This work was supported by the New York Marathon Limb Preservation Fund.

REFERENCES

- Amudsen MA, Spittel JA, Thompson JH, Owen CA, Jr: Hypercoagulability associated with malignant disease and with the postoperative state. *Ann Intern Med* 1963;58:608-616.
- Clagett GP, Reisch JS: Prevention of venous thromboembolism in general surgical patients. *Ann Surg* 1988;208:227-240.
- Edwards EA: Migrating thrombophlebitis associated with carcinoma. *N Engl J Med* 1949;240:1031-1035.
- Menzioan JO, Sequeira JC, Doyle JE, et al.: Therapeutic and clinical course of deep vein thrombosis. *Am J Surg* 1983;146:581-585.
- Rickles FR, Levine M, Edwards RL: Hemostatic alterations in cancer patients. *Cancer Metastasis Rev* 1992;1:237-248.
- Sproul EF: Carcinoma and venous thrombosis: The frequency of association of carcinoma in the body or tail of the pancreas with multiple venous thrombosis. *Am J Cancer* 1938;34:566-585.
- Byrd RB, Divertie MB, Spittel JA, Jr: Bronchogenic carcinoma and thromboembolic disease. *J Am Med Assn* 1967;202:1019-1022.
- Clarke-Pearson DL, Coleman RE, Synan IS, et al.: Venous thromboembolism prophylaxis in gynecologic oncology: A prospective controlled trial of low-dose heparin. *Am J Obstet Gynecol* 1983;145:606-613.
- Clarke-Pearson DL, Synan IS, Dodge R, et al.: A randomized trial of low-dose heparin and intermittent pneumatic calf compression for the prevention of deep venous thrombosis after gynecologic oncology surgery. *Am J Obstet Gynecol* 1993;168:1146-1153.
- Coe NP, Collins REC, Klein LA, et al.: Prevention of deep vein thrombosis in urological patients: A controlled randomized trial of low dose heparin and external pneumatic compression boots. *Surgery* 1978; 83:230-234.
- Cubilla A, Fitzgerald PJ: Pancreas cancer. I. Duct adenocarcinoma: A clinical-pathologic study of 380 patients. *Pathol Annual* 1978;13:241-289.
- Goodnough LT, Saito H, Manni A, et al.: Increased incidence of thromboembolism in stage IV breast cancer patients treated with a five drug chemotherapy regimen: A study of 159 patients. *Cancer* 1984;54:1264-1268.
- Pinzon R, Drewinko B, Trujillo JM, et al.: Pancreatic carcinoma and Trousseau's syndrome: Experience at a large cancer center. *J Clin Oncol* 1986;4:509-514.
- Sons HU, Borchard F: Esophageal cancer autopsy findings in 171 cases. *Arch Pathol Lab Med* 1984;108:983-988.
- Sons HU, Borchard F, Joel BS: Carcinoma of the gallbladder autopsy findings in 287 cases and review of the literature. *J Surg Oncol* 1985;28:199-206.
- Haake DA, Berkman SA: Venous thromboembolic disease after hip surgery. Risk factors, prophylaxis, and diagnosis. *Clin Orthop* 1989;242:212-231.
- Johnson R, Green JR, Charnley J: Pulmonary embolism and its prophylaxis following the Charnley total hip replacement. *Clin Orthop* 1977;127:123-132.
- Lieberman JR, Geerts WH: Prevention of venous thromboembolism after total hip and knee arthroplasty. *J Bone Joint Surg* 1994;76-A:1239-1250.
- Evarts CM, Feil EJ: Prevention of thromboembolic disease after elective surgery of the hip. *J Bone Joint Surg* 1971;53-A:1271-1280.
- Harris WH, Salzman EW, Athanasoulis CA, et al.: Aspirin prophylaxis of venous thromboembolism after total hip replacement. *N Engl J Med* 1977;297:1246-1249.
- Rogers PH, Walsh PN, Marder VJ, et al.: Controlled trial of low-dose heparin and sulfipyrazone to prevent venous thromboembolism after operation on the hip. *J Bone Joint Surg* 1978;60-A:758-762.
- Sagar S, Nairn D, Stamatakis JD, et al.: Efficacy of low-dose heparin in prevention of extensive deep vein thrombosis in patients undergoing total-hip replacement. *Lancet* 1976;1:1151-1154.
- Sikorski JM, Hampson WG, Staddon GE: The natural history and aetiology of deep vein thrombosis after total hip replacement. *J Bone Joint Surg* 1981;63-B:171-177.
- Haas SB, Tribus CB, Insall JN, et al.: The significance of calf thrombi after total knee arthroplasty. *J Bone Joint Surg* 1992;74-B:799-802.
- Leclerc JR, Geerts WH, Desjardins L, et al.: Prevention of deep vein thrombosis after major knee surgery: A randomized, double-blind trial comparing a low molecular weight heparin fragment (enoxaparin) to placebo. *Thromb Haem* 1992;67:417-423.
- Lynch JA, Baker PL, Polly RE, et al.: Mechanical measures in the prophylaxis of postoperative thromboembolism in total knee arthroplasty. *Clin Orthop* 1990;260:24-29.
- McKenna R, Bachmann F, Kaushal SP, Galante JO: Thromboembolic disease in patients undergoing total knee replacement. *J Bone Joint Surg* 1976;58-A:928-932.
- Stringer MD, Steadman CA, Hedges AR, et al.: Deep vein thrombosis after elective knee surgery. An incidence study in 312 patients. *J Bone Joint Surg* 1989;71-B:492-497.
- Stulberg BN, Insall JN, Williams GW, Ghelman B: Deep-vein thrombosis following total knee replacement: An analysis of six hundred and thirty-eight arthroplasties. *J Bone Joint Surg* 1984;66-A:194-201.
- Balderston RA, Graham TS, Booth RE, Rothman RH: The prevention of pulmonary embolism in total hip arthroplasty: Evaluation of low-dose warfarin therapy. *J Arthroplasty* 1989;3:217-221.
- Harris WH, McCusick K, Athanasoulis CA, et al.: Detection of pulmonary emboli after total hip replacement using serial C¹⁵O₂ pulmonary scans. *J Bone Joint Surg* 1984;66-A:1388-1393.
- Lotke PA, Steinberg ME, Ecker ML: Significance of deep venous thrombosis in the lower extremity after total joint arthroplasty. 1994; *Clin Orthop* 299:25-30.
- Wolf LD, Hozack WJ, Rothman RH: Pulmonary embolism in total joint arthroplasty. *Clin Orthop* 1993;288:219-233.
- Haas SB, Insall JN, Scuderi GR, et al.: Pneumatic sequential-compression boots compared with aspirin prophylaxis of deep-vein thrombosis after total knee arthroplasty. *J Bone Joint Surg* 1990;72-A:27-31.
- Lynch AF, Bourne RB, Rorabeck CH, et al.: Deep-vein thrombosis and continuous passive motion after total knee arthroplasty. *J Bone Joint Surg* 1988;70-A:11-14.
- Culver D, Crawford JS, Gardiner JH, Wiley AM: Venous thrombosis after fractures of the upper end of the femur: A study of incidence and site. *J Bone Joint Surg* 1970;52-B:61-69.
- Geerts WH, Code KI, Jay RM, et al.: A prospective study of venous thromboembolism after major trauma. *N Engl J Med* 1994; 331:1601-1606.
- Hamilton HW, Crawford JS, Gardiner JH, Wiley AM: Venous thrombosis in patients with fracture of the upper end of the femur. A phlebographic study of the effect of prophylactic anticoagulation. *J Bone Joint Surg* 1970;52-B:268-289.
- Montgomery KD, Geerts WH, Potter HG, Helfet DL: Thromboembolic complications in patients with pelvic trauma. *Clin Orthop* 1996;329:68-87.
- Moskovitz PA, Ellenberg SS, Feffer HL, et al.: Low-dose heparin for prevention of venous thromboembolism in total hip arthroplasty and surgical repair of hip fractures. *J Bone Joint Surg* 1978;60-A:1065-1070.
- Sharnoff JG, Rosen RL, Sadler AH, Ibarra-Isunza GC: Prevention of fatal pulmonary thromboembolism by heparin prophylaxis after surgery for hip fractures. *J Bone Joint Surg* 1976;58-A:913918.
- Svend-Hansen H, Bremerskov V, Gotrik J, Osti P: Low-dose heparin in proximal femoral fractures: Failure to prevent deep-vein thrombosis. *Acta Orthop Scandinavica* 1081;52:77-80.
- Davis RB, Theologides A, Kennedy BJ: Comparative studies of blood coagulation and platelet aggregation in patients with cancer and nonmalignant disease. *Ann Int Med* 1969;71:67-80.
- Rickles FR, Hancock WW, Edwards RL, Zacharski LR: Antimetastatic agents: I. Role of cellular procoagulants in the pathogenesis of fibrin deposition in cancer and the use of anticoagulants and/or

- antiplatelet drugs in cancer treatment. *Sem Thromb Hem* 1988;14:88-94.
45. Reston VA: ACR standards for performance of the peripheral venous ultrasound examination. Res 6 1993.(Abstract) AUTHOR: Please provide name of meeting.
 46. Cronan JJ, Dorfman GS, Grusmark J: Lower-extremity deep venous thrombosis: Further experience with and refinements of US assessment. *Radiology* 1988;168:101-107.
 47. Hartman JT, Pugh JL, Smith RD, et al.: Cyclic sequential compression of the lower limb in prevention of deep venous thrombosis. *J Bone Joint Surg* 1982; 64-A:1059-1062.
 48. Hull R, Delman TJ, Hirsch J, et al.: Effectiveness of intermittent pulsatile elastic stockings for the prevention of calf and thigh vein thrombosis in patients undergoing elective knee surgery. *Thromb Res* 1979;16:37-45.
 49. Hull RD, Raskob GE, Gent M, et al.: Effectiveness of intermittent pneumatic leg compression for preventing deep vein thrombosis after total hip replacement. *JAMA* 1990;263:2313-2317.
 50. Paiement G, Wessinger SJ, Waltman AC, Harris WH: Low dose warfarin versus external pneumatic compression for prophylaxis against venous thromboembolism following total hip replacement. *J Arthroplasty* 1987;2:23-26.
 51. Woolson ST, Watt JM: Intermittent pneumatic compression to prevent proximal deep venous thrombosis during and after total hip replacement: A prospective, randomized study of compression alone, compression and aspirin, and compression and low-dose warfarin. *J Bone Joint Surg* 1991;73-A:507-512.
 52. Allenby F, Boardman L, Pflug JJ, Calman JS: Effects of external pneumatic intermittent compression on fibrinolysis in man. *Lancet* 1973;2:1412-1414.
 53. Knight MT, Dawson R: Effect of intermittent compression of the arms on deep venous thrombosis in the legs. *Lancet* 1976;2:1265-1267.
 54. Weitz J, Michelsen J, Gold K, et al.: Effects of intermittent pneumatic calf compression in post-operative thrombi and plasmin activity. *Thromb Haem* 1986;56:198-201.
 55. Bailey JP, Kruger MP, Solano FS, et al.: Prospective randomized trial of sequential compression devices vs low-dose warfarin for deep venous thrombosis prophylaxis in total hip arthroplasty. *J Arthroplasty* 1991;6:29-35.
 56. Francis CW, Pelligrini Jr. VD, Marder VJ, et al.: Comparison of warfarin and external pneumatic compression in prevention of venous thrombosis after total hip replacement. *JAMA* 1992;267:2911-2915.
 57. Mohr DN, Silverstein MD, Murtaugh PA, Harrison JM: Prophylactic agents for venous thrombosis in elective hip surgery: Meta-analysis of studies using venographic assessment. *Arch Intern Med* 1993;153:2221-2228.
 58. Oishi CS, Grady-Benson JC, Otis SM, et al.: The clinical course of distal deep venous thrombosis after total hip and total knee arthroplasty as determined with duplex ultrasonography. *J Bone Joint Surg* 1994;76-A:1658-1663.
 59. Aitken AGF, Godden DJ: Real-time ultrasound diagnosis of deep vein thrombosis: A comparison with venography. *Clin Radiol* 1987;38:309-313.
 60. Appelman PT, De Jong TE, Lampmann LE: Deep venous thrombosis of the leg: US findings. *Radiology* 1987;163:743-746.
 61. Dauzat MM, Laroche JP, Charras C, et al.: Real-time B-mode ultrasonography for better specificity in the noninvasive diagnosis of deep venous thrombosis. *J Ultrasound Med* 1986;5:625-631.
 62. Raghavendra BN, Rosen RJ, Lam S, et al.: Deep venous thrombosis. Detection by high-resolution real-time ultrasound. *Radiology* 1984;152:789-793.
 63. Froehlich JA, Dorfman GS, Cronan JJ, et al.: Compression ultrasonography for the detection of deep venous thrombosis in patients who have a fracture of the hip: A prospective study. *J Bone Joint Surg* 1989;71-A:249-256.
 64. Grady-Benson JC, Oishi CS, Hanson PB, et al.: Postoperative surveillance for deep venous thrombosis with duplex ultrasonography after total knee arthroplasty. *J Bone Joint Surg* 1994;76-A:1649-1657.
 65. Wells PS, Lensing AW, Davidson BL, et al.: Accuracy of ultrasound for the diagnosis of deep venous thrombosis in asymptomatic patients after orthopedic surgery: A meta-analysis. *Ann Intern Med* 1995;122:47-53.
 66. Woolson ST, McCrory DW, Walter JF, et al.: B-mode ultrasound scanning in the detection of proximal venous thrombosis after total hip replacement. *J Bone Joint Surg* 1990;72-A:983-987.
 67. Leutz DW, Stauffer ES: Color duplex doppler ultrasound scanning for detection of deep venous thrombosis in total knee and hip arthroplasty patients: Incidence, location, and diagnostic accuracy compared with ascending venography. *J Arthroplasty* 1994;9:543-548.
 68. Amstutz HC, Friscia DA, Dorey F, Carney BT: Warfarin prophylaxis to prevent mortality from pulmonary embolism after total hip replacement. *J Bone Joint Surg* 1989;71-A:321-326.
 69. Colwell CW Jr., Spiro TE, Trowbridge AA, et al.: Use of enoxaparin, a low-molecular-weight heparin and unfractionated heparin for the prevention of deep venous thrombosis after elective hip replacement. *J Bone Joint Surg* 1994;76-A:3-14.
 70. Guyer RD, Booth RE, Rothman RH: The detection and prevention of pulmonary embolism in total hip replacement: A study comparing aspirin and low-dose warfarin. *J Bone Joint Surg* 1982;64-A:1040-1044.
 71. Harris WH, Salzman EW, Athanasoulis C, et al.: Comparison of warfarin, low-molecular-weight dextran, aspirin, and subcutaneous heparin in prevention of venous thromboembolism following total hip replacement. *J Bone Joint Surg* 1974;56-A:1552-1562.
 72. Hull R, Raskob G, Pineo G, et al.: A comparison of subcutaneous low-molecular-weight heparin with warfarin sodium for prophylaxis against deep-vein thrombosis after hip or knee implantation. *N Engl J Med* 1993;329:1370-1376.
 73. Nurmohamed MT, Rosendaal FR, Buller HR, et al.: Low-molecular-weight heparin versus standard heparin in general and orthopaedic surgery: A meta-analysis. *Lancet* 1992;340:152-156.
 74. Paiement GD, Wessinger SJ, Hughes R, Harris WH: Routine use of adjusted low-dose warfarin to prevent venous thromboembolism after total hip replacement. *J Bone Joint Surg* 1993;75-A:893-898.
 75. Turpie AG, Levine MN, Hirsh J, et al.: A randomized controlled trial of a low-molecular-weight heparin (enoxaparin) to prevent deep-vein thrombosis in patients undergoing elective hip surgery. *N Engl J Med* 1986;315:925-929.
 76. Vresilovic Jr. EJ, Hozack WJ, Booth RE, Rothman RH: Incidence of pulmonary embolism after total knee arthroplasty with low-dose coumadin prophylaxis. *Clin Orthop* 1993;286:27-31.
 77. Bick RL: Coagulation abnormalities in malignancy: A review. *Sem Thromb Hem* 1992;18:353-372.
 78. Krauth D, Holden A, Knapic N, et al.: Safety and efficacy of long-term oral anticoagulation in cancer patients. *Cancer* 1987;59:983-985.